

Cu requirement for periodic drawing of blood from the animals (Deignan, T., et al., Res Vet Sci 69(2) 153-58 (2000)). The amount of IgG (antibody) obtained is usually between 3-8 mg/ ml of serum. The method also involves bleeding of the rabbit several times to obtain the antibodies, as the titer is highest only between the 8th - 10th day after 2 - 3 boosters. Monoclonal antibodies are produced by immunizing an animal with a protein, obtaining antibody-producing cells from the animal, and fusing the antibody-producing cells with strains of myeloma cells, i.e., tumor cells, to produce hybridomas, which are isolated and cultured as monoclones. The monoclonal hybridomas may either be cultured in vitro or from the cells, ascitic fluid, or serum of a tumor-bearing host animal. Since each antibody-producing cell generates a single unique antibody, each monoclonal culture of hybridomas produces a homogeneous antibody. Not all of the hybridoma clones, which result from fusing myeloma cells with antibody-producing cells, are specific for the desired pesticide (of functional groups upon the pesticide characteristic of that class of molecules), since many of the hybridomas will make antibodies, which the inoculated animal has produced to react with other foreign substances. Even antibodies against the subject antigen will differ from clone to clone, since antibodies produced by different cells may react with different antigenic determinants of the same molecule. From each clone, therefore, it is necessary to obtain the resulting antibody and test its reactivity with the subject pesticides and to test its specificity by determining which particular organochlorine pesticide it recognizes. Further, only certain antibodies or antisera function in specific immunoassay formats or configurations.

Page 2, second paragraph:

92 United States Patent Nos. 4,357,272 (Polson, A.) and 4,550,019 (Polson, A.) claimed production of hen egg yolk antibodies and have been used in a number of applications for passive transfer of immunity.

Page 3, last paragraph:

GA Reference is made to the animal studies of Yolken, R. et al., Pediatrics 81(2): 291-295 (1988) and Yolken, R. et al., Journal Clin. Immunology 10(6Suppl): 80S-87S (1990), proposed the oral administration of antiviral HEY immunoglobulin for the prevention and treatment of enteric infections, including rotaviral infection in humans. Methods and formulations for the oral administration of immune globulin are known (U.S. Pat No. 4,477,432 to Hardie). However, there are no reports on the production of egg yolk antibodies for a small molecule such as a pesticide.

Applicants respectfully request that the foregoing amendments be made prior to examination of the present application.

Respectfully submitted,

Date January 24, 2003

By



FOLEY & LARDNER
Washington Harbour
3000 K Street, N.W., Suite 500
Washington, D.C. 20007-5109
Telephone: (202) 672-5300
Facsimile: (202) 672-5399

Matthew E. Mulkeen
Registration No. 44,250

Should additional fees be necessary in connection with the filing of this paper, or if a petition for extension of time is required for timely acceptance of same, the Commissioner is hereby authorized to charge Deposit Account No. 19-0741 for any such fees; and applicant(s) hereby petition for any needed extension of time.